

Andrew Biggs*, Daniel Zalduendo**

* The Vale Veterinary Group, The Laurels, Station Road, Tiverton Devon EX164LF UK, Tel:- +44 (0) 1884 259584 Email:- ValeLab@btinternet.com

** HIPRA, Avda. La Selva, n°135 17170 – Amer (Girona) Spain

1. Objective: Apply mastitis diagnostic and monitoring techniques to three herds chosen for a high prevalence of *Staphylococcus aureus* over the 12 months prior to, and during the 12 months of, a rolling 3 month vaccination program with a polyvalent mastitis vaccine. STARTVAC® (HIPRA).

2. Vaccination protocol

- All cows received 2 vaccine doses 3 week apart followed by a rolling policy of quarterly boosters.
- In-calf heifers were batched to receive 2 doses 3 weeks apart with the second dose no less than 10 days prior to expected parturition. Heifers then joined the rolling 3 month booster program.

3. Data observations:

- Data is compared during 6 month intervals from 12 months prior to and 12 months after initiating a whole herd STARTVAC® vaccination policy for:

- Clinical cases
- *S. aureus* prevalence
- Somatic Cell Count

↑ = Start of vaccination

4. Conclusions from data to date

In general:

- Better results after 6 months using the rolling protocol - in common with other studies the improvements take time.
- Vaccination is not a panacea and consistent good mastitis management is essential.
 - Herd De had some issues with both the parlour and environment resulting in increased challenge elevating the new infection rate.
- Clinical mastitis reduced by 54%
- *S. aureus* prevalence reduced by 71%
- Somatic Cell Count
 - Percentage of healthy animals increased by 4%
 - First infections reduced by 49%

